



TITLE:

Moduli Spaces of Holomorphic Mappings into Hyperbolically Imbedded Complex Spaces and Locally Symmetric Spaces(Automorphic Forms and Related Topics)

AUTHOR(S):

Noguchi, Junjiro

CITATION:

Noguchi, Junjiro. Moduli Spaces of Holomorphic Mappings into Hyperbolically Imbedded Complex Spaces and Locally Symmetric Spaces(Automorphic Forms and Related Topics). 数理解析研究所講究録 1987, 617: 218-230

ISSUE DATE:

1987-03

URL:

<http://hdl.handle.net/2433/99837>

RIGHT:

**Moduli Spaces of Holomorphic Mappings into Hyperbolically
Imbedded Complex Spaces and Locally Symmetric Spaces**

Junjiro Noguchi

野口 潤次郎

Department of Mathematics, Faculty of Science
Tokyo Institute of Technology
Oh-Okayama, Meguroku, Tokyo 152, Japan

Introduction

The aim of this talk is to describe the results of Noguchi [14]. Let X be a connected Zariski open subset of a compact reduced complex space \bar{X} such that X is complete hyperbolic and hyperbolically imbedded into \bar{X} (cf. [7, 16]). Let N be a Zariski open subset of a compact complex manifold \bar{N} such that $\partial N = \bar{N} - N$ is a hypersurface with only normal crossings; in some case, we consider the case of $\partial N = \emptyset$. Here we study the structure of the moduli space $\text{Hol}(N, X)$ of all holomorphic mappings $f: N \rightarrow X$ of N into X . Especially interesting is the case where X is the quotient $\Gamma \backslash D$ of a

symmetric bounded domain D by a torsion-free arithmetic subgroup Γ of the identity component $\text{Aut}^0(D)$ of the holomorphic transformation group $\text{Aut}(D)$ of D . It is known that $\Gamma \backslash D$ is complete hyperbolic and hyperbolically imbedded into the Satake compactification $\overline{\Gamma \backslash D}$ of $\Gamma \backslash D$ (cf. [10, 2, 8, 9]). Besides the interesting results of [20, 21, 18], the present work is motivated by the results on the Parshin-Arakelov theorems for curves [17, 1] and Abelian varieties [5]. Cf. also [13]. Let $\pi: \bar{Y} \rightarrow \bar{N}$ be a fiber space over \bar{N} which is smooth over N , such that the fibers $Y_x = \pi^{-1}(x)$ with $x \in N$ are curves with a given genus g or g -dimensional Abelian varieties with principal polarization. Then, roughly speaking, the fiber space naturally induces a holomorphic mapping $f: N \rightarrow \Gamma \backslash S_g$, where S_g denotes Siegel's generalized upper half space. Then the deformation of $\pi: \bar{Y} \rightarrow \bar{N}$ as fiber space over \bar{N} with degeneration at most over ∂N and the total space of such fiber spaces correspond respectively to the deformation of the holomorphic mapping f and the moduli space $\text{Hol}(N, \Gamma \backslash S_g)$. Thus it is quite natural to deal with the case where N and $\Gamma \backslash D$ are non-compact. In the case where N is compact, there is an earlier work for a fiber space of Abelian varieties by [11].

§1. Holomorphic mappings into hyperbolically imbedded spaces

The natural topology of $\text{Hol}(N, X)$ which we endow with is the compact-open topology. We first prove an extension

and convergence theorem.

Theorem (1.1). Let X be a hyperbolic complex space and hyperbolically imbedded into \bar{X} . Let N be a complex manifold \bar{N} minus a hypersurface with only normal crossings. If a sequence $\{f_\nu\}_{\nu=1}^\infty$ of $f_\nu \in \text{Hol}(N, X)$ converges to a holomorphic mapping $f: N \rightarrow \bar{X}$, then there are unique holomorphic extensions $\bar{f}_\nu: \bar{N} \rightarrow \bar{X}$ of f_ν and $\bar{f}: \bar{N} \rightarrow \bar{X}$ of f , and $\{\bar{f}_\nu\}$ converges uniformly on compact subsets of \bar{N} to \bar{f} .

As for the extension theorem, this generalizes the result of [7], but the method of the proof is different. It will play a fundamental role in our arguments. In the proof of Theorem (1.1) we use the following lemma (cf. [14] for the details).

Lemma (1.2) (cf. [19]). Let $B(R)$ be the open ball of the m -dimensional complex vector space \mathbb{C}^m with radius R and center 0 . Let S be an analytic subset of pure dimension k of $B(R)$ such that $0 \in S$. Then we have

$$\text{Vol}(S \cap B(r)) \geq \frac{\pi^k}{k!} r^{2k}$$

for $0 < r < R$. Moreover, if the equality holds for some $r > 0$, then S is a linear subspace of \mathbb{C}^m .

In what follows, we assume that \bar{N} and \bar{X} are compact, and that X is a Zariski open subset of \bar{X} , complete hyperbolic and hyperbolically imbedded into \bar{X} . Combining Theorem (1.1) with the Douady theory [3], we have

Theorem (1.3). Hol(N, X) carries a structure of a complex space with universal property, such that its underlying topology coincides with the compact-open topology, and

$$\Phi: (f, x) \in \text{Hol}(N, X) \times N \rightarrow f(x) \in X$$

is a holomorphic mapping, which is proper for every fixed $x \in N$. Moreover, $\text{Hol}(N, X)$ is a Zariski open subset of a compact complex space.

Sketch of the proof. Let $\text{Hol}(\bar{N}, \bar{X})$ be the space of all holomorphic mappings from \bar{N} into \bar{X} with compact-open topology. Then, by Theorem (1.1) the mapping

$$f \in \text{Hol}(N, X) \rightarrow \bar{f} \in \text{Hol}(\bar{N}, \bar{X})$$

is an into-homeomorphism. Hence we identify the topological space $\text{Hol}(N, X)$ with its image in $\text{Hol}(\bar{N}, \bar{X})$. By making use of the distance decreasing property of hyperbolic distance for holomorphic mappings, we see that $\text{Hol}(N, X)$ is relatively compact. The complete hyperbolicity of X implies that $\text{Hol}(N, X)$ is open and closed in $\text{Hol}(\bar{N}, \bar{X})$ and then Theorem (1.1) yields that the topological closure of $\text{Hol}(N, X)$ in $\text{Hol}(\bar{N}, \bar{X})$ is a compact complex subspace which contains $\text{Hol}(N, X)$ as a Zariski open subset. The complete hyperbolicity of X also implies that $\Phi(\cdot, x): \text{Hol}(N, X) \rightarrow X$ is proper for every fixed $x \in N$. Q.E.D.

In general, let Y_1 and Y_2 be two complex spaces. For a holomorphic mapping $f: Y_1 \rightarrow Y_2$, we set

$$\text{rank } f = \sup \left\{ \dim_t Y_1 - \dim_t f^{-1}(f(t)); t \in Y_1 \right\}.$$

The following proposition follows from Lemma (1.2). It reveals a special nature of the complex analyticity of holomorphic mappings but is less known.

Proposition (1.4). Assume that Y_1 and Y_2 are compact. Let $\{f_\nu\}_{\nu=1}^\infty$ be a sequence of points of $\text{Hol}(Y_1, Y_2)$ converging to $f \in \text{Hol}(Y_1, Y_2)$. If $\text{rank } f_\nu = k$, then $\text{rank } f = k$.

We set

$$\text{Hol}(k; N, X) = \{f \in \text{Hol}(N, X); \text{rank } f = k\}.$$

Corollary (1.5). $\text{Hol}(k; N, X)$ is open and closed in $\text{Hol}(N, X)$.

§2. The moduli $\text{Hol}(N, \Gamma \backslash D)$

In this section we deal with the case where X is the quotient $\Gamma \backslash D$ of a symmetric bounded domain D by a torsion-free discrete subgroup Γ of $\text{Aut}(D)$. We assume that Γ is uniform or an arithmetic subgroup of $\text{Aut}^0(D)$. In the case where Γ is uniform and $N = \bar{N}$, the results of this section were already obtained in [20, 21, 18]. We are mainly interested in the case where $\Gamma \backslash D$ and N are non-compact, while our arguments work in the compact case. Let $l(D)$ (resp. $l(\Gamma)$) denote the maximum dimension of proper boundary components of D (resp. Γ -rational boundary components). Let

$\text{Hol}(k; N, \Gamma \backslash D)$ denote the set of all holomorphic mappings $f: N \rightarrow \Gamma \backslash D$ with $\text{rank } f = k$. Applying the results of the previous section, we have

Theorem (2.1). i) $\text{Hol}(N, \Gamma \backslash D)$ carries a structure of a complex space compatible with compact-open topology, such that the evaluation mapping

$$\Phi: (f, x) \in \text{Hol}(N, \Gamma \backslash D) \times N \rightarrow f(x) \in \Gamma \backslash D$$

is holomorphic. Moreover, $\text{Hol}(N, \Gamma \backslash D)$ is a Zariski open subset of the compact complex space $\overline{\text{Hol}(N, \Gamma \backslash D)}^{1)}$, and satisfies the universality property; i.e., for a complex space T and a holomorphic mapping $\psi: T \times N \rightarrow \Gamma \backslash D$, the natural mapping

$$t \in T \rightarrow \psi(t, \cdot) \in \text{Hol}(N, \Gamma \backslash D)$$

is holomorphic.

ii) Every connected component of $\text{Hol}(N, \Gamma \backslash D)$ is complete hyperbolic and the holomorphic mappings

$$\Phi_x: f \in \text{Hol}(N, \Gamma \backslash D) \rightarrow f(x) \in \Gamma \backslash D$$

are proper for all $x \in N$.

1) $\overline{\text{Hol}(N, \Gamma \backslash D)}$ is the closure of $\text{Hol}(N, \Gamma \backslash D)$ in $\text{Hol}(\bar{N}, \Gamma \backslash D)$.

- iii) $\text{Hol}(k; N, \Gamma \setminus D)$ are open and closed in $\text{Hol}(N, \Gamma \setminus D)$.
- iv) $\text{Hol}(k; N, \Gamma \setminus D)$ are compact for $k > \ell(\Gamma)$.
- v) $\text{Hol}(k; N, \Gamma \setminus D)$ are finite for $k > \ell(D)$.

In the last of this section, we study in details the structure of $\text{Hol}(N, \Gamma \setminus D)$, assuming that \bar{N} is Kähler and ∂N is a hypersurface with only simple normal crossings. We use the following result on harmonic mappings by [18]:

(2.2) Let $F: N \rightarrow \Gamma \setminus D$ and $G: N \rightarrow \Gamma \setminus D$ be free homotopic harmonic mappings with finite energy. Then there is a harmonic mapping $\Psi: \mathbb{R} \times N \rightarrow \Gamma \setminus D$ with respect to the product metric $dt \oplus h$ on $\mathbb{R} \times N$ such that

- i) $\Psi(0, x) = F(x)$, $\Psi(1, x) = G(x)$ and Ψ provides a free homotopy between F and G , equivalent to the given one;
- ii) for every $x \in N$, the curve $\gamma_x: t \in \mathbb{R} \rightarrow \Psi(t, x) \in \Gamma \setminus D$ is a parametrization of a geodesic with constant speed, independent of x , and $e(\Psi(t, \cdot))(x)$ is constant in t .

Lemma (2.3). Let F and G be as in (2.2). If F is holomorphic, then so is G .

Remark. 1) In case N is compact, this is a theorem due to Lichnerowicz (cf. Theorem (8.6) of [4]).

2) Since (2.2) actually holds for harmonic mappings from a complete Riemannian manifold with finite volume into a complete Riemannian manifold with non-positive sectional curvatures, Lemma (2.3) is also true for harmonic mappings F and G from a complete Kähler manifold with finite volume into a complete Kähler manifold with non-positive sectional curvatures, provided that F and G have finite energies.

The main result is the following:

Theorem (2.4). i) $\text{Hol}(N, \Gamma \backslash D)$ is smooth and quasi-projective.

ii) For a connected component Z of $\text{Hol}(N, \Gamma \backslash D)$ and a point $x \in N$, the evaluation mapping at x

$$\phi_x: f \in Z \rightarrow f(x) \in \Gamma \backslash D$$

is a proper holomorphic immersion onto a totally geodesic complex submanifold, so that Z is a free quotient of a symmetric bounded domain.

iii) For a connected component Z of $\text{Hol}(N, \Gamma \backslash D)$, there is a normal complex projective variety \tilde{Z} such that Z is hyperbolically imbedded into \tilde{Z} and ϕ_x holomorphically extends to $\bar{\phi}_x: \tilde{Z} \rightarrow \overline{\Gamma \backslash D}$.

iv) $\dim \text{Hol}(k; N, \Gamma \backslash D) \leq \ell(D)$ for $k > 0$.

- v) For $f \in \text{Hol}(N, \Gamma \backslash D)$ with $\bar{f}^{-1}(\partial \Gamma \backslash D) \neq \emptyset$,
 $\dim_f \text{Hol}(N, \Gamma \backslash D) \leq \ell(\Gamma)$.

As a corollary, we have the following.

Corollary (2.5) (Rigidity). Let $f: N \rightarrow \Gamma \backslash D$ be a holomorphic mapping. Then f is a unique holomorphic mapping among the free homotopy class of f , if f satisfies one of the following conditions:

- a) The image of f is not contained in a totally geodesic complex proper submanifold of $\Gamma \backslash D$;
- b) $\text{rank } f > \ell(D)$;
- c) $\bar{f}^{-1}(\partial \Gamma \backslash D) \neq \emptyset$ and $\text{rank } f > \ell(\Gamma)$.

In general, a holomorphic mapping $f \in \text{Hol}(N, \Gamma \backslash D)$ admits a deformation (cf. [5]). But in the special case where D is the n -th product H^n of the upper half plane $H \subset \mathbb{C}$, we see that any $f \in \text{Hol}(N, \Gamma \backslash H^n)$ is rigid. That is, by making use of the rigidity Theorem 6 of [6], we have

Theorem (2.6). Let $\Gamma \subset (\text{PSL}(2, \mathbb{R}))^n$ be an irreducible torsion-free discrete subgroup with $\text{Vol}(\Gamma \backslash H^n) < \infty$. Then

- i) if $f: N \rightarrow \Gamma \backslash H^n$ is a non-constant holomorphic mapping, f is a unique holomorphic mapping among the free homotopy class of f ,

so that

ii) there are only finitely many non-constant holomorphic mappings from N into $\Gamma \backslash H^n$.

Remark. 1) It must be noted that if $\Gamma \backslash H^n$ is not compact, then Γ is arithmetic ([12]). Therefore Γ satisfies our requirement for discrete subgroups.

2) In the case of $\dim N = 1$, i) was proved in [6].

3) By the same arguments as in [15], we see that the Kähler assumption for \bar{N} is not necessary in ii). The proof is reduced to the present case.

4) For a compact quotient $\Gamma \backslash H^2$ and a compact complex manifold N , ii) was proved in [15]. For an algebraic curve N and compact $\Gamma \backslash H^n$, it was proved in [6].

References

- [1] S. Ju. Arakelov, Families of algebraic curves with fixed degeneracies, Izv. Akad. Nauk SSSR ser. Mat., 35 (1971), 1277-1302.
- [2] A. Borel, Some metric properties of arithmetic quotients of symmetric spaces and an extension theorem, J. Differential Geometry, 6 (1972), 543-560.

- [3] A. Douady, Le problème des modules pour les sous-espaces analytiques compacts d'un espace analytique donné, *Ann. Inst. Fourier, Grenoble*, 16 (1966), 1-95.
- [4] J. Eells and L. Lemaire, Selected topics in harmonic maps, *CBMS*, 50, Amer. Math. Soc., 1983.
- [5] G. Faltings, Arakelov's theorem for Abelian varieties, *Invent. Math.*, 73 (1983), 337-347.
- [6] Y. Iwayoshi, Generalizations of de Franchis theorem, *Duke Math. J.*, 50 (1983), 393-408.
- [7] P. Kiernan, Extensions of holomorphic maps, *Trans. Amer. Math. Soc.*, 172 (1972), 347-355.
- [8] P. Kiernan, On the compactifications of arithmetic quotients of symmetric spaces, *Bull. Amer. Math. Soc.*, 80 (1974), 109-110.
- [9] P. Kiernan and S. Kobayashi, Comments on Satake compactification and the great Picard theorem, *J. Math. Soc. Japan*, 28 (1976), 577-580.
- [10] S. Kobayashi and T. Ochiai, Satake compactification and the great Picard theorem, *J. Math. Soc. Japan*, 23 (1971), 340-350.
- [11] M. Kuga and S.-I. Ihara, Family of families of Abelian varieties, *Algebraic Number Theory*, Kyoto, 1976, Japan Soc. for the Promotion of Science, Tokyo, 1977, 126-

142.

- [12] G. A. Margulis, Arithmeticity of non-uniform lattices, Functional Anal. Appl., 7 (1973), 245-246.
- [13] D. Mumford, Curves and their Jacobians, Univ. of Michigan Press, Ann Arbor, 1975.
- [14] J. Noguchi, Moduli spaces of holomorphic mappings into hyperbolically imbedded complex spaces and locally symmetric spaces, preprint.
- [15] J. Noguchi and T. Sunada, Finiteness of the family of rational and meromorphic mappings into algebraic varieties, Amer. J. Math., 104 (1982), 887-900..
- [16] T. Ochiai and J. Noguchi, Geometric Function Theory in Several Complex Variables, Iwanami, Tokyo, 1984. (in Japanese)
- [17] A. N. Parshin, Algebraic curves over function fields. I, Izv. Akad. Nauk SSSR Ser. Mat., 32 (1968), 1145-1170.
- [18] R. Schoen and S. T. Yau, Compact group actions and the topology of manifolds with non-positive curvature, Topology, 18 (1979), 361-380.
- [19] G. Stolzenberg, Volumes, Limits, and Extensions of Analytic Varieties, Lecture Notes in Math., 19, Springer-Verlag, Berlin-Heidelberg-New York, 1966.

[20] T. Sunada, Holomorphic mappings into a compact quotient of symmetric bounded domain, Nagoya Math. J., 64 (1976), 159-175.

[21] T. Sunada, Rigidity of certain harmonic mappings, Invent. Math., 51 (1979), 297-307.